

Crowdsourcing gas flux climatologies and the big experiment

Jamie Shutler, Jean-Francois Piolle, Peter Land, David Woolf, Fanny Girard-Ardhuin,
Bertrand Chapron and Craig Donlon



PML | Plymouth Marine
Laboratory



UNIVERSITY OF
EXETER



oceanflux ghg
support to science element



Crowdsourcing gas flux climatologies and the big experiment

- 30 people took part.
- 39 individual processing runs.
- 6 failures (5 users gave no email address, 1 actual failure)
- 8 completed, but with no data (ASCAT data option not available)



Crowdsourcing gas flux climatologies and the big experiment

- 30 people took part.
- 39 individual processing runs.
- 6 failures (5 users gave no email address, 1 actual failure)
- 8 completed, but with no data (ASCAT data option not available)
- 25 completed runs:
 - 17 valid runs, Mean global net sink = $-1.7 \pm 0.51 \text{ PgC yr}^{-1}$




Crowdsourcing gas flux climatologies and the big experiment

- 30 people took part.
- 39 individual processing runs.
- 6 failures (5 users gave no email address, 1 actual failure)
- 8 completed, but with no data (ASCAT data option not available)
- 25 completed runs:
 - 17 valid runs, Mean global net sink = $-1.7 \pm 0.51 \text{ PgC yr}^{-1}$
 - 5 added biases to k, mean net sink = 86 PgC yr^{-1} !
 - 2 rain k runs
 - 1 produced nans !



Crowdsourcing gas flux climatologies and the big experiment



Participant preferences:

- 80% used Takahashi, 20% used SOCATv2
- 72% used non-bulk flux formulation
- 60% used backscatter based gas transfer k

